arXiv:1411.3570v1 [math.MG] 13 Nov 2014

PROXIMAL VORONOÏ REGIONS

J.F. PETERS

Dedicated to the Memory of Som Naimpally

Abstract. A main result in this paper is the proof that proximal Voronoï regions are convex polygons. In addition, it is proved that every collection of proximal Voronoï regions has a Leader uniform topology.

1. Introduction

Klee-Phelps convexity [8, 12] and related results [11] are viewed here in terms of Voronoï regions, named after the Ukrainian mathematician Georgy Voronoï [13, 14, 15]. A nonempty set A of a space X is a *convex set*, provided $\alpha A + (1-\alpha)A \subset A$ for each $\alpha \in [0,1]$ [1, §1.1, p. 4]. A *simple convex set* is a closed half plane (all points on or on one side of a line in R^2).

Lemma 1.1. [6, §2.1, p. 9] *The intersection of convex sets is convex.*

Proof. Let $A, B \subset \mathbb{R}^2$ be convex sets and let $K = A \cap B$. For every pair points $x, y \in K$, the line segment $\overline{xy}$ connecting x and y belongs to K, since this property holds for all points in A and B. Hence, K is convex. □



Figure 1. V_p = Intersection of closed half-planes

Let $S \subset \mathbb{R}^2$ be a finite set of n points called sites, $p \in S$. The set S is called the *generating set* [7]. Let H_{pq} be the closed half plane of points at least as close to p as to $q \in S \setminus \{p\}$, defined by

$$H_{pq} = \left\{ x \in R^2 : \|x - p\| \underset{q \in S}{\leq} \|x - q\| \right\}.$$

2010 *Mathematics Subject Classification.* Primary 65D18; Secondary 54E05, 52C20, 52C22.

Key words and phrases. Convex polygon, proximal, Leader uniform topology, Voronoï region.

The research has been supported by the Scientific and Technological Research Council of Turkey (TÜBİTAK) Scientific Human Resources Development (BIDEB) under grant no: 2221-1059B211402463 and Natural Sciences & Engineering Research Council of Canada (NSERC) discovery grant 185986.

A *convex polygon* is the intersection of finitely many half-planes [5, §I.1, p. 2]. See, for example, Fig. 1.

Remark 1.2. The Voronoï region V_p depicted as the intersection of finitely many closed half planes in Fig. 1 is a variation of the representation of a Voronoï region in the monograph by H. Edelsbrunner [6, §2.1, p. 10], where each half plane is defined by its outward directed normal vector. The rays from p and perpendicular to the sides of V_p are comparable to the lines leading from the center of the convex polygon in G.L. Dirichlet's drawing [3, §3, p. 216]. ■

2. Preliminaries

Let $S \subset E$, a finite-dimensional normed linear space. Elements of S are called sites to distinguish them from other points in E [6, §2.2, p. 10]. Let $p \in S$. A *Voronoï region* of $p \in S$ (denoted V_p) is defined by

$$V_p = \left\{ x \in E : \|x - p\| \underset{\forall q \in S}{\leq} \|x - q\| \right\}.$$

Remark 2.1. A Voronoï region of a site $p \in S$ contains every point in the plane that is closer to p than to any other site in S [7, §1.1, p. 99]. Let V_p, V_q be Voronoï polygons. If $V_p \cap V_q$ is a line, ray or line segment, then it is called a *Voronoï edge*. If the intersection of three or more Voronoï regions is a point, that point is called a *Voronoï vertex*. ■

Lemma 2.2. *A Voronoï region of a point is the intersection of closed half planes and each region is a convex polygon.*

Proof. From the definition of a closed half-plane

$$H_{pq} = \left\{ x \in R^2 : \|x - p\| \underset{q \in S}{\leq} \|x - q\| \right\},$$

V_p is the intersection of closed half-planes H_{pq}, for all $q \in S - \{p\}$ [5], forming a polygon. From Lemma 1.1, V_p is a convex. □

A Voronoi diagram of S (denoted by $\mathbb{V}$) is the set of Voronoi regions, one for each site $p \in S$, defined by

$$\mathbb{V} = \bigcup_{p \in S} V_p.$$

Example 2.3. Centroids as Sites in an Image Tessellation.
Let E be a segmentation of a digital image and let $S \subset E$ be a set of sites, where each site is the centroid of a segment in E. In a centroidal approach to the Voronoï tessellation of E, a Voronoï region V_p is defined by the intersection of closed half plains determined by centroid $p \in S$. The centroidal approach to Voronoi tessellation was introduced by Q. Du, V. Faber, M. Gunzburger [4]. ■

3. Main Results

Let V_p, V_z be Voronoï regions of $p, z \in S$, a set of Voronoï sites in a finite-dimensional normed linear Space E that is topological, cl$A$ the closure of a nonempty set A in E. V_p, V_z are *proximal* (denoted by $V_p \ \delta \ V_z$), provided $\mathbb{P} = \text{cl}V_p \cap \text{cl}V_z \neq \emptyset$ [2]. The set $\mathbb{P}$ is called a *proximal Voronoï region*.

Theorem 3.1. *Proximal Voronoï regions are convex polygons.*

Proof. Let $\mathbb{P}$ be a proximal Voronoï region. By definition, $\mathbb{P}$ is the nonempty intersection of convex sets. From Lemma 1.1, $\mathbb{P}$ is convex. Consequently, $\mathbb{P}$ is the intersection of finitely many closed half planes. Hence, from Lemma 2.2, $\mathbb{P}$ is a Voronoï region of a point and is a convex polygon. □

Corollary 3.2. *The intersection of proximal Voronoï regions is either a Voronoï edge or Voronoï point.*

Any two Voronoï regions intersect at least a vertex and at most along their boundaries. Together, the set of Voronoï regions $\mathbb{V}$ cover the entire plane [5, §2.2, p. 10]. For a set of sites $S \subset E$, a Voronoï diagram $\mathbb{D}$ of S is the set of Voronoi regions, one for each site in S.

Corollary 3.3. *A Voronoï diagram $\mathbb{D}$ equals $\mathbb{V}$.*

The partition of a plane E with a finite set of n sites into n Voronoï polygons is known as a Dirichlet tessellation, named after G.L. Dirichlet [16] (see [3]). A *cover* (covering) of a space X is a collection $\mathcal{U}$ of subsets of X whose union contains X (*i.e.*, $\mathcal{U} \supseteq X$) [17, §15], [10, §7.1].

Corollary 3.4. *A Dirichlet tessellation $\mathbb{D}$ of the Euclidean plane E is a covering of E.*

Recall that the Euclidean space $E = R^2$ is a metric space. The topology in a metric space results from determining which points are close to each set in the space. A point $x \in E$ is close to $A \subset E$, provided the Hausdorff distance $d(x, A) = inf\{\|x - a\| : a \in A\} = 0$. Let X, Y be a pair of metric spaces, $f : X \longrightarrow Y$ is a function such that for each $x \in X$, there is a unique $f(x) \in Y$. A continuous function preserves the closeness (proximity) between points and sets, *i.e.*, $f(x)$ is close to $f(B)$ whenever x is close to B. In a proximity space, one set A is near another set B, provided $A\ \delta\ B$, *i.e.*, the closure of A has at least one element in common with the closure of B. The set A is close to the set B, provided the Čech distance $D(A, B) = inf\{\|a - b\| : a \in A, b \in B\} = 0$. In that case, we write $A\ \delta\ B$ (A and B are proximal). A *uniformly continuous mapping* is a function that preserves proximity between sets, *i.e.*, $f(A)\ \delta\ f(B)$ whenever $A\ \delta\ B$. A *Leader uniform topology* is determined by finding those points that are close to each given set in E.

Theorem 3.5. *Let S be a set of two or more sites, $p \in S, V_p \in \mathbb{D}$ in the Euclidean space R^2. Then*
1º V_p is near at least one other Voronoï region in $\mathbb{D}$.
2º Let p, y be sites in S. $\{y\}\ \delta\ \{p\} \Rightarrow \{y\}\ \delta\ V_p$.
3º V_p is close to Voronoï region V_y if and only if $d(x, V_y) = 0$ for at least one $x \in V_p$.
4º A mapping $f : V_p \longrightarrow V_y$ is uniformly continuous, provided $f(V_p)\ \delta\ f(V_y)$ whenever $V_p\ \delta\ V_y$.

Proof.
1º: Assume S contains at least 2 sites. Let $p \in S, y \in S \setminus \{y\}$ such that V_p, V_y have at least one closed half plane in common. Then $V_p\ \delta\ V_y$.
2º: If $\{y\}\ \delta\ \{p\}$, then $\|y - p\| = 0$, since $y \in \{y\} \cap \{p\}$. Consequently, $\{y\} \cap \text{cl}(V_p) \neq \emptyset$. Hence, $\{y\}\ \delta\ \text{cl}(V_p)$.
3º: $V_p\ \delta\ V_y \Leftrightarrow$ exists $x \in \text{cl}(V_p) \cap \text{cl}(V_y) \Leftrightarrow d(x, V_y) = 0$.

4^o: Let $f(V_p)\ \delta\ f(V_y)$ whenever $V_p\ \delta\ V_y$. Then, by definition, $f : V_p \longrightarrow V_y$ is uniformly continuous. □

Theorem 3.6. *Every collection of proximal Voronoï regions has a Leader uniform topology (application of* [9]*).*

Proof. Assume $\mathbb{D}$ has more than one Voronoï region. For each $V_p \in \mathbb{D}$, find all $V_y \in \mathbb{D}$ that are close to V_p. For each V_p, this procedure determines a family of Voronoï regions that are near V_p. Let τ be a collection of families of proximal Voronoï regions. Let $A, B \in \tau$. $A \cap B \in \tau$, since either $A \cap B = \emptyset$ or, from Theorem 3.5.l^o, there is at least one Voronoï region $V_p \in A \cap B$, *i.e.*, $V_p\ \delta\ A$ and $V_p\ \delta\ B$. Hence, $A \cap B \in \tau$. Similarly, $A \cup B \in \tau$, since $V_p\ \delta\ A$ or $V_p\ \delta\ B$ for each $V_p \in A \cup B$. Also, $\mathbb{D}, \emptyset$ are in τ. Then, τ is a Leader uniform topology in $\mathbb{D}$. □

References

[1] G. Beer, *Topologies on Closed and Closed Convex Sets*, Kluwer Academic Pub.,Boston, MA, 1993, MR1269778.

[2] A. Di Concilio, *Proximity: A powerful tool in extension theory, function spaces, hyperspaces, Boolean algebras and point-free geometry*, Amer. Math. Soc. Contemporary Math. **486** (2009), 89114, MR2521943.

[3] G.L. Dirichlet, *Über die Reduktion der positiven quadratischen Formen mit drei unbestimmten ganzen Zahlen*, Journal für die reine und angewandte **40** (1850), 221-239.

[4] Q. Du, V. Faber and M. Gunzburger, *Centroidal Voronoï tessellations: Applications and algorithms*, SIAM Review **41** (1999), no. 4, 637-676, MR1722997.

[5] H. Edelsbrunner, *Geometry and topology of mesh generation*, Cambridge University Press, Cambridge, UK, 2001, 209 pp., MR1833977.

[6] H. Edelsbrunner, *A Short Course in Computational Geometry and Topology*, Springer, Berlin, 110 pp.

[7] N.P. Frank and S.M. Hart, *A Dynamical System Using the Voronoï Tessellation*, The Amer. Math. Monthly **117** (2010), no. 2, 99112, MRMR2590195.

[8] V.L. KLee, *A characterization of convex sets*, The Amer. Math. Monthly **56** (1949), no. 4, 247249, MR0029519.

[9] S. Leader, *On clusters in proximity spaces*, Fundamenta Mathematicae **47** (1959), 205213, MRMR0112120.

[10] S.A. Naimpally and J.F. Peters, *Topology with applications. Topological spaces via near and far*, World Scientific, Singapore, 2013, xv + 277pp, MR3075111, Zbl 1295.68010.

[11] J.F. Peters, M.A. Öztürk and M. Uçkun, *Klee-Phelps convex groupoids*, arXiv **1411.0934v1 [math.GR]** (2014), 1-4.

[12] R.R. Phelps, *Convex sets and nearest points*, Proc. Amer. Math. Soc. **8** (1957), no. 4, 790-797, MR0087897.

[13] G. Voronoï, *Sur un problème du calcul des fonctions asymptotiques*, J. für die reine und angewandte **126** (1903), 241-282, JFM 38.0261.01.

[14] G. Voronoï, *Nouvelles applications des paramètres continus à la théorie des formes quadratiques. Premier Mémoir*, J. für die reine und angewandte **133** (1908), 97-178, JFM 38.0261.01.

[15] G. Voronoï, *Nouvelles applications des paramètres continus à la théorie des formes quadratiques. Deuxième Mémoir*, J. für die reine und angewandte **134** (1908), 198-287, JFM 39.0274.01.

[16] E.W. Weisstein, *Voronoi Diagram*, Wolfram MathWorld (2014), http://mathworld.wolfram.com/VoronoiDiagram.html.

[17] S. Willard, *General Topology*, Addison-Wesley Pub. Co., Reading, Mass., 1970, xii + 369 pp., MR0264581.

E-mail address: James.Peters3@umanitoba.ca

Computational Intelligence Laboratory, University of Manitoba, WPG, MB, R3T 5V6, Canada